



Multiple Sclerosis Modulating the Epigenome

Dr. Syed Mohamed Ibrahim Sulthan
(Ph.D., from Delhi University & Working at Shaqra University)
Bio-Medical Engineering Laboratory,
College of Engineering, Shaqra University,
Ad-Dawadimi, Riyadh Province-11911, SAUDI ARABIA

Abstract: Multiple sclerosis (MS) may be an advanced malady characterised by autoimmune degenerative disorder of the medullary sheath of the central systema nervosum. Variable degrees of incapacity result from interruptions in psychological, physical, and neural phenomena. No clear reason for MS has been known, however specific genetic mutations and environmental effects are established as factors in illness development. However, the low sickness concordance rates and transmission disequilibrium determined in MS forestall genetic factors alone from explaining magnified susceptibleness. As a result, the involvement of epigenetics, or transmissible changes within the ordination that aren't caused by alterations within the underlying desoxyribonucleic acid sequence, has been projected to resolve these apparent discrepancies. This paper focuses on recent analysis into 3 primary epigenetic mechanisms and their relevance to MS susceptibility: Desoxyribonucleic acid methylation, simple protein modifications, and post-transcriptional regulation of target genes by microRNA. These epigenetic phenomena act together with one another and environmental stimuli to provide malady.

Keywords: degenerative disorder; protein modifications; methylation; genetic mutations; microRNA

I. Introduction

Multiple sclerosis (MS) is that the most typical demyelinating sickness of the central systema nervosum, touching 2.0 to 2.5 million individuals worldwide [37] [41]. MS defies conventional classification, possessing each autoimmune and neurologic attributes, and is characterized through the development of demyelinated foci within the central systema nervosum and by a range of chronic physical, neurologic, and psychological symptoms [29] [30] [62].

Demyelination, or injury to the sheath that surrounds axons of the brain and medulla spinalis, is that the results of focal infiltration with the aid of antibodies, macrophages and, lymphocytes which results in brief irritation [25] [34]. Efforts at remyelination do occur, however usually the repair of those lesions isn't permanent and consequent remyelination efforts are progressively more unsuccessful [12]. The physical and neurologic disabilities that characterize MS are the results of the accretion of injury over time [57].

MS has constantly been ascertained all through a larger percentage of the population in international locations aloof from the equator. North American country specially has one in all uppermost rates of MS worldwide, collectively with a numeral of countries in Europe (northern region) [60]. Charles Poser recommended that maybe MS was propagated genetically through Scandinavian lineages, as MS happens with such high frequency in Scandinavian countries. Abundant proof supports this idea of MS vulnerability, with factors like climate, month of birth, and sun exposure acting as environmental stimuli that influence sickness hazard and development through mediation by genetic and epigenetic mechanisms [29] [30] [62]. Alternative factors, like smoking and EBV infection, contribute additional to influence an individual's vulnerability to MS [44].

II. Multiple Sclerosis and Epigenetics

Changes within the desoxyribonucleic acid sequence of MS-related genes are solely a minor aspect in the advancement of the malady. Budding knowledge recommend that epigenetics, or changes in genetic appearance that aren't a consequence of changes within the underlying desoxyribonucleic acid sequence, might have a higher influence on the advancement and progression of MS than mutations in any specific locus [5] [29] [30] [62]. Epigenetic marks are sustained through generations of somatic cells among the body, further as in gametes, and might thus be passed from blood relation to kids [5] [27]. However, as a result of their elevated reaction to environmental situational therapy, epigenetic marks are distinctively proficient of being altered or inverted over time [27]. 3 principal epigenetically-controlled mechanisms participate key functions within the

pathophysiology of MS: deoxyribonucleic acid methylation, histone modifications, and microRNA-associated post-transcriptional cistron regulation.

A. DNA methylation

DNA methylation is presently the chief most extensively analysed epigenetic technique, and engross the addition of a methyl cluster group to cytosine residues in CpG dinucleotides among a deoxyribonucleic acid sequence via deoxyribonucleic acid-methyltransferase (DNMT) action [34]. Hypermethylated cytosine filtrates inside a promoter section intrude with sequence identification through transcription elements. As an end result, this kind of amendment is normally associated with the steady repression of gene expression at the related locus [24] [31].

B. Histone Variations

Genes may also be regulated epigenetically via modifications to the superstructure of DNA. One such aspect of this condensed shape is histones, acute phase protein just about which DNA is twisting [14]. Gene expression can be modulated by using numerous post-translational modifications, which include methylation or acetylation, and the combined effects of those changes confer a unique histone code to first-rate cell types [13] [23]. the general mechanism via which gene expression is altered thru modifications in chromatin shape that have an effect on the accessibility of unique loci genetic to transcription machinery [13] [29] [30]. Whether or not genetic appearance is repressed or activated relies upon the actual mixture of adjustments which might be discovered on histones, with a view to both condense the chromatin to repress expression or increase it to improve transcription [1].

C. miRNA-associated put up-transcription

MicroRNAs (miRNAs) are tiny fiber, non-coding molecules fashioned from larger transcripts that fold to supply pin structures [53]. When many nuclear and cytoplasmatic process steps, miRNAs complicated with a bunch of proteins to supply the RNA- precipitated silencing complicated (RISC) that allows miRNA to alter gene expression by binding to related messenger RNA transcripts [63]. Imperfect base pairing reasons the target messenger RNA to be degraded, whereas wonderful base pairing induces amendment of location repression of the target's sequence product via the inhibition of translation [63].

III. The Role of Epigenetics in MS

The clinical attributes of MS recommend that underlying epigenetic mechanisms might contribute to malady receptiveness and progression. For example, the low concordance occurrence of MS in twins (roughly five percent in dizygous twins and 25-30% in monozygotic twins) means that genetic polymorphisms alone can't be the reason of the sickness considering that identical twins proportions one hundred percent of their genes [12] [51]. Additionally, epigenetic mechanisms are thought to mediate genomic acquisition which will account for the enhanced prevalence of MS in ladies. For example, the DRB1*15 allelomorph of the HLA genes, that is concerned in immunity and is related to an elevated MS risk, is carried by a lot of ladies than men [8].

Although the epigenetic mechanisms represented here have their own distinct effects, it's necessary to notice that these processes intermingle instead of act in isolation. Cancer studies recommend that deoxyribonucleic acid methylation and histamine cephalgia modifications are capable of modulating miRNA expression, that themselves are regulated by alternative miRNAs [19] [29] [30]. Alternative interactions occur between alkyl group deoxyribonucleic acid and histone-modifying enzymes, like accomplishment of DNMTs by histones to induce de novo deoxyribonucleic acid methylation [6]. These primary epigenetic mechanisms moreover exert their outcomes together with specific environmental factors [29] [30] [62].

Taken along, diverse components of MS pathophysiology and sickness development could also be produced as a results of epigenetic adjustments that set off outcomes induce of their personal, at the same time as additionally interacting in complicated networks that are not yet absolutely understood.

A. DNA methylation

Methylation of CpG dinucleotides in some of gene promoter areas that can be answerable for the immune characteristics of MS have been described. The immunopathology of the disorder entails the dominance of T helper cell 1 (TH1) immunity, related to the cytokine interferon- γ (IFN- γ), over TH2 cell immunity [32]. The differentiation of the 2 T cell sorts is regulated through epigenetic mechanisms, as is the production of IFN- γ and different cytokines [32]. Aberrant patterns of DNA methylation were discovered inside the promoter location for IFN- γ inside T helper cells, which may be associated with the dominance of TH1 immunity over TH2 immunity in MS [32].

Analyses of brain tissue samples from MS sufferers have identified an affiliation between DNA methylation and neuronal cellular loss of life [10]. Here we have a look at, overexpression of the enzyme answerable for de novo methylation of DNA triggered apoptosis in cells. Furthermore, expression of other DNMTs enhanced while apoptosis become precipitated.

Smoking is thought to be a hazard aspect for mounting MS and is related to more frequent relapses and more numerous MS lesions in sufferers [2]. Commonly thru DNA hypermethylation, research on the link among

cigarette smoking and most cancers have determined that smoking induces the silencing of tumour suppressor genes [58].

Hypermethylation of the promoter areas of tumour suppression genes is likewise associated with Epstein-Barr virus (EBV) contamination [43]. Viral latent membrane protein 1 acts to up-modify diverse DNMTs, which in turn induce hypermethylation at diverse promoter locations, articulated from research regarding most cancers [56]. In fact, individuals with a history of infective mononucleosis, a kind of EBV contamination, double their hazard of growing MS due to adjustments in the epigenetic profile of inflamed cells [20].

Hypomethylation of some promoters also can have an effect on MS improvement. For example, while the promoter of the PAD2 gene is hypomethylated, and the overexpression of this enzyme is observed [36]. Citrullination of myelin basic protein (MBP) by PAD2, and this is a key participant in MS pathophysiology. The changed form of MBP is much less barn, and the formation of neutral citrullinyl residues from positively-charged arginyl residues might bring about much less compact myelin this is prone to disintegration and a probable autoimmune reaction to MBP [36] [40]. Proven white substance that appeared standard for MS patients brain biopsy samples, however contained extended degrees of citrullinated MBP in assessment to samples from healthful control youngsters, adults, and individuals with different neurodegenerative sicknesses, consisting of Alzheimer's, Huntington's or Parkinson's [38].

B. Histone modifications and MS pathogenesis

Some of the modifications to histones were found to be associated with the development and demonstration of MS. In a study of sufferers with progressive MS and controls without neurological disorder, an boom in histone H3 acetylation (the addition of acetyl clusters) in neural cells referred to as oligodendrocytes within chronic MS lesions was discovered, whereas early- phase MS lesions contained oligodendrocytes with marked histone deacetylation (the elimination of acetyl groups) [48]. The researchers cited that in rodent fashions, histone deacetylation materialize to support differentiation of oligodendrocytes, while acetylation ended in impaired differentiation because of enhanced expression of transcriptional inhibitors of cellular differentiation [48]. Diminished oligodendrocyte differentiation might result in impaired remyelination, imparting a capability link to lesion improvement [29] [30]. Moreover, the analysis established that histone H3 acetylation is improved in samples from older sufferers and the quantity of acetylation is correlated with the severity of sickness[48].

There is a huge effect of histone structure on oligodendrocyte improvement. Histone deacetylases (HDACs: enzymes that dispose of acetyl clusters from histone tails) inhibit development into unconventional cell lineages and repress myelin-unique gene inhibitors in oligodendrocyte relatively undifferentiated cells [54]. HDAC1 is recruited to the proximal promoter vicinity of MBP genes by means of transcription factors Nkx2.2 and Hes5 [59]. Those changes in gene expression mediate oligodendrocyte maturation, a procedure which is regularly aberrant in MS and this ends up in impaired remyelination [13].

HDAC inhibitors had been shown to have tremendous effects in animal models of MS. Camelo et al. located in 2005 that administering trichostatin A (TSA) to mice with experimental autoimmune encephalomyelitis (EAE), an animal model of MS, attenuated spinal twine infection, demyelination and axonal loss. Prevention of acylation of histones, blocking the IL-12 signalling pathway in T cells and preventing EAE from growing in mice, are possible by a selective simple protein acetyltransferase inhibitor; according to Natajara et al. No similar studies had been undertaken in people, however these observations lend help to the function of histone changes in MS and other demyelinating problems.

Biochemical alterations to histones play a task inside the extended risk that vitamin D deficiency confers to MS. In reality, epidemiological studies advise that as blood levels of 25-hydroxyvitamin D3 increase, the chance of growing MS reduces [39]. Individuals with relapse-remitting MS generally tend to experience extra incapacity and extra frequent relapses with decreased stages of vitamin D, that is complex by the truth that they will experience difficulty in obtaining solar exposure, a chief factor in vitamin D production, due to physical incapacity [55]. The way in which vitamin D exerts an effect on MS susceptibility and severity is doubtful, however studies on nutrition D deficiency and most cancers propose that it can epigenetically impact ailment by using interacting with histones: 25-hydroxyvitamin D induces the expression of JMJD3, a gene that encodes histone H3 demethylase, an enzyme that removes methyl clusters from histones, and thereby alters the overall chromatin shape of goal genes [49]. Vitamin D is similarly connected with sturdy immune-mediating capabilities, and is even capable of stopping the onset and development of EAE in mice [7]. Mice that have been administered nutrition D on the onset of neurological signs, then later had this treatment discontinued, resumed EAE development [7].

The impact of nutrition D on MS is likewise implicated inside the extended incidence of MS in people born in the spring than the ones born in disdain months, as concluded by means of Willer and colleagues in their assessment of MS patients in Canada, along with comparable statistics pooled from Britain, Denmark, Sweden, and other international locations [60]. Those investigations indicated that a drastically higher wide variety of patients with MS had been born in May than those born in November, and that this modification may be related to the higher geographical latitude of the nations studied and for that reason the extent of solar coverage in those areas [60]. It has now not been conclusively hooked up why these styles of improved MS chance were

discovered, and in addition research into these findings might be an interesting and worthwhile region of providence analysis.

In the end, histone citrullination, which is the enzymatic conversion of arginine residues to citrulline residues, has a big effect at the etiopathogenesis of MS as fighting fit. In the presence of augmented stages of tumour necrosis aspect α (TNF α), the enzyme PAD4, which is related to PAD2, relocates from the cytoplasm to the nucleus and additionally enhances histone citrullination [13] [35]. Determined that citrullination occurs prior to the onset of signs and symptoms in mice with EAE and located an enhance in histone citrullination in the white substance of MS patients in evaluation to control themes.

C. miRNA-associated post-transcriptional regulation in MS

The majority of our contemporary information of how MS is regulated by way of epigenetic mechanisms is derived from research that concentrates on sufferers with actively-demyelinating lesions. In a single analysis, awesome miRNA profiles had been determined in biopsy samples from active MS lesions from affected people that showed tremendous up-regulation of miR-155, miR-34a, and miR-326 [28]. The miRNA transcripts of this family of genes intention CD47, a cell surface protein that inhibits the phagocytic commotion of macrophages and is broadly expressed in cells [28]. The miRNAs decrease CD47 expression, which results in the activation of macrophages and subsequently, the phagocytic destruction of myelin, for this reason producing an active lesion if the cell is not able to result in remyelination [9] [28].

Some other cluster of research observed that miR-155 is likewise related to T cell differentiation, with miR-326 appearing similar features [46]. Up-regulation of miR-one hundred fifty five in a few kinds of leukocytes (macrophages, B cells, and T cells), in reaction to ligand fastening to toll-like receptors and inflammation cytokines, indicates that miRNA can be involved in an inflammatory reaction [46]. It has additionally been discovered that mice deficient in miR-155 are resistant to growing EAE when in comparison to wild-type mice [46]. Similarly, an analysis of MS sufferers and mice with EAE discovered that miR-326 was exceptionally up-regulated in peripheral blood cells, and this growth in expression turned into most reported in the course of patient relapses, indicating that miR-326 is pretty correlated with sickness severity [15]. In comparison, control subjects and MS sufferers in remission exhibited reduced levels of miR-326 expression [15]. miR-326 aims a transcription factor that inhibits T cell differentiation; consequently, overexpression of miR-326 suppresses the expression of this protein and ultimately will enhance manufacturing of persuaded varieties of T cells [15]. This occasion can result in the immune adjustments related to MS improvement.

Up-regulation of other miRNA genes has been implicated in MS. Similarly to miR-one hundred fifty five, extended expression of miR-338 and miR-491 in mind samples, acquired from patients with revolutionary MS, prompted the transcribed miRNAs to target and suppress the translation of enzymes inside the aldo-keto reductase own family, that are involved within the production of neurosteroids, which include allopregalone, in brain white subject matter [45]. Researchers determined that in mice with EAE, overexpression of miR-338 and miR-491 decreased the production of allopregalone, whereas mice that were administered the neurosteroid exhibited decreased brain infection and attenuated myelin and axonal harm, suggesting that allopregalone possesses neuroprotective characteristics[45]. Endogenous neurosteroid synthesis is regulated through miRNA-associated put up-transcriptional silencing, and as such, the dysregulation of synthesis is an important pathological mechanism in revolutionary MS [45].

IV. Expectations in Epigenetics of MS

As epigenetics is a pretty new vicinity of studies, maximum studies of the epigenetic mechanisms concerned in MS are preliminary, but open doors for future studies that could extrapolate from the ideas they present and explore prospective applications in extra detail. They'll additionally provide to clarify discrepancies among findings in studies of similar phenomena: as an example, not all proof has supported DNA methylation as a sizeable issue in MS improvement. Baranzini et al. undertook research assessing genetic, epigenetic, and transcriptomic variations in CD4⁺ T helper cells between three monozygotic dual pairs, wherein one twin suffered from MS, in a challenge and give an explanation for malady discordance. But, no applicable variations in DNA methylation styles had been found inside the genomes of T cells in co-twins, offering little help for explaining discordance, and suggesting that other epigenetic mechanisms can also play a role in MS pathogenesis [3]. Inside the identical year, Ramagopalan and Ebers cautioned that because somatic cells collect and regulate their epigenetic marks over the direction of a person's lifetime in reaction to their environment, monozygotic twins won't, in fact, have same gene expression, and therefore might have substantial differences in their epigenome. They affirm, but, that although the Baranzini et al. research seems to tentatively rule out epigenetic changes as a issue in MS development, other cell varieties need to be examined and the genomes of a number twin pairs must be sequenced for you to make greater correct judgments about the influence of epigenetics in sickness improvement [51]. Furthermore, because the charges of sequencing research, which include the one undertaken through Baranzini et al., proceed to reduce, new revelations may also provide stepping stones for important future research guidelines.

Different popular guidelines may be made for prospective studies in MS epigenetics. Larger study populations have to be employed in research involving each fashionable MS sufferers and twins. Research on twins and disorder concordance quotes which have been published so far have used very small cohorts of just a few dual pairs, and therefore may not constitute intensive information of the way epigenetics in MS is involved in concordance and heredity. Moreover, present research, particularly latest publications, must be replicated to validate present day findings, preferably with large cohorts.

Of specific hobby is analysis which hyperlinks neuroimaging with epigenetic findings, magnetic resonance imaging (MRI) and positron emission tomography in particular useful. Evaluation might be designed that evaluate specific epigenetic marks located in MS sufferers with lesion improvement through the years in mind concern or with the kind of MS exhibited through the affected person. In addition, longitudinal studies of MS sufferers and the epigenetic marks they appear would be useful. Researchers may want to have a look at changes in DNA methylation, purposeful agencies connected to histone tails, or miRNA expression at a couple of intervals throughout a person's lifetime, and examine whether or not there are variations through the years. Are the epigenetic marks solid? Do they correlate with disease severity or periods of remission? Family unit with a couple of affected participants are also beneficial take a look at topics as they offer the opportunity to investigate whether epigenetic marks related to MS improvement are propagated stably from dad and mom to their children.

A few researches are finished that look into using DNA methylation as a biomarker for sickness activity in most cancer [18] [33]. Extending the ones programs to MS might offer a reliable manner that of diagnosing the malady, and possibly in predicting the path of relapse and remission. Sodium bisulphite mapping could be a helpful experimental device for deciding the unique CpG islands inside a gene that are methylated. Research concerning this method is also accustomed compile a library of methylation sites that are of best importance in MS patients.

Upregulation of DNMTs in necrobiosis may be concerned within the neuronc degeneration observed in MS, however no research have analysed their upregulation as a potential disorder mechanism. It should be prudent to start this research in destiny studies of EAE in animal models, probably mice, with EAE as an illustration of MS. Mastronardi et al. projected an experimental therapeutic approach that epigenetically reprograms oligodendrocytes with the aid of favouring methylation on the promoter of IFN- γ genes. INF- β is presently a general remedy in relapse-remitting MS for decreasing the range of symptomatic episodes, however combinatorial remedy of IFN- β with nutrition B12 down-regulated IFN- γ expression in *in vitro* studies of oligodendrocytes [64]. IFN- β alone extended the quantity of strategies in growing oligodendrocytes, however combinatorial remedy significantly elevated oligodendrocyte maturation, additionally as attenuated infection [64]. Due to the fact *in vitro* experiments created effective ends up in oligodendrocyte maturation, *in vivo* research regarding this combinatorial therapy are attainable next steps.

The analysis performed by Pedre et al. instructed that histone acylation and different histone modifications could additionally have a function within the pathologic process and physiology of MS, however that similarly studies is essential so as to extend our understanding of the results of those modifications to histones. The final functional outcomes of histone adjustments ought to be tested as nicely. A exquisite deal of analysis has began on the jobs that miRNA play within the development of MS by way of interacting with the genome, and future analysis may additionally uncover different miRNAs that have an effect on MS pathologic process. it can be useful to inquire as to whether or not transcription factors or different tiny molecules additionally manage gene expression in the disease. when you consider that an appropriate functions of every miRNA could also be troublesome to decide in people because of the multitude of different factors that contribute to MS pathophysiology, victimisation of unique cellular traces or animal fashions can be greater feasible.

There are different queries that stay that relate to MS susceptibleness and ailment development. Even though it has been broadly stated that EBV contamination will increase an individual's MS hazard, it remains to be confirmed specifically how this improved chance is generated. is that the increase associated to the changes within the epigenetic profile of infected cells as recommended with the aid of Handel and co-workers? What kinds of epigenetic changes in those cells are related with MS? More analysis is needed to work out the position of Epstein-Barr virus in MS pathologic process and the way it conjures autoimmunity in affected people.

MS researchers maintain to create strides in extra elucidating illness mechanisms and growing novel treatments, enhancements with the intention to truly contribute to our collective statistics of MS and therefore the remaining well-being of MS sufferers. The examination of epigenetics in MS is growing considerably our angle at the complex factors that engage to supply the disorder.

V. Conclusion

We've got mentioned molecular mechanisms of the control of gene expression that occur independently of modifications within the desoxyribonucleic acid collection. These epigenetic mechanisms embrace post-translational adjustments of nucleosomal histones, desoxyribonucleic acid methylation, and regulation by way

of non-coding miRNAs, and may be laid low with the environment and lifestyle (eg, weight loss program, diet, smoking behavior, exercising, and drug addiction). We advise that those environmental factors would possibly regulate sickness manifestation by using modulating the epigenome, which multiple sclerosis would possibly result from accumulative adjustments imposed by environmental factors at the immune system and within the brain. Awesome cell populations would possibly differentially answer to the equal stimulus, by changing the association of the nuclear structure during a cell- unique manner. In immune cells, epigenetic control would possibly result in extended expression of genes related to sickness manifestation, whereas within the brain, similar stimuli would possibly impair the flexibility of progenitors to shape new myelin. As an instance, social interplay turned into recently proven to profoundly have an effect on myelin content within the prefrontal cortex of adult mice, and to modulate put up-translational adjustments of nucleosomal histones. The mixture of epigenetic changes in immune cells and oligodendrocytes would possibly modulate illness onset (by using modulating expression of susceptibleness genes) and ailment course (through modulating myeline formation). But, warning is guaranteed concerning use of epigenetic marks as sickness biomarkers for neurological problems, due to the fact that epigenetic adjustments diagnosed in a specific tissue population (eg, marginal blood monocytes) cannot be effortlessly generalised to a exclusive organ (eg, mind) or cell kind (eg, CD4+ lymphocytes). Despite the fact that, among a single tissue, epigenetic mechanisms want to be studied longitudinally to work out the result of environmental stimuli on disorder direction or healing responsiveness. The presence of restrictive marks in areas containing sequences specific for nutrition D receptor binding, for instance, would give an explanation for the dearth of patient responsiveness to nutrition D supplementation. It may be anticipated that maps of responsive genes are going to be created and used for screening sufferers within the future. This may be viable handiest if specific signatures of epigenetic maps are appropriately outlined for every cell type. The combination of the wealth of knowledge from genome-huge association research with the longitudinal analysis of epigenetic marks on chosen patient populations will probably result in effective individualized healing approaches.

References

- [1] Alva V, Ammelburg M, Soding J, Lupas A. 2007. On the origin of the histone fold. *BMC Structural Biology*. 7:17.
- [2] Ascherio A, Munger KL. 2007. Environmental risk factors for multiple sclerosis. Part II: noninfectious factors. *Annals of Neurology*. 61:504-513.
- [3] Baranzini SE, Mudge J, van Velkinburgh JC, Khankhanian P, Khrebtukova I, Miller NA, Zhang L, Farmer AD, Bell CJ, Kim RW, et al. 2010. Genome, epigenome and RNA sequences of monozygotic twins discordant for multiple sclerosis. *Nature*. 464:1351-1356.
- [4] Bartel DP. 2004. MicroRNAs: genomics, biogenesis, mechanism and function. *Cell*. 116:281-297.
- [5] Berger SL, Kouzarides T, Shiekhattar R, Shilatifard S. 2009. An operational definition of epigenetics. *Genes and Development*. 23:781-783.
- [6] Camelo S, Iglesias AH, Hwang D, Due B, Ryu H, Smith K, Gray SG, Imitola J, Duran G, Assaf B et al. 2005. Transcriptional therapy with the histone deacetylase inhibitor trichostatin A ameliorates experimental autoimmune encephalomyelitis. *Journal of Immunology*. 164(1-2):10-21.
- [7] Cantorna MT, Hayes CE, Deluca HF. 1996. 1,25-dihydroxyvitamin D3 reversibly blocks the progression of relapsing encephalomyelitis, a model of multiple sclerosis. *Proceedings of the National Academy of Sciences*. 93:7861-7864.
- [8] Chao M.J, Herrera BM, Ramagopalan SV, Deluca G, Handunnetthi L, Orton SM, Lincoln MR, Sadovnick AD, Ebers GE. 2010. Parent-of-origin effects at the major histocompatibility complex in multiple sclerosis. *Human Molecular Genetics*. 19:3679-3689.
- [9] Chari D. 2007. Remyelination in multiple sclerosis. *International Review of Neurobiology*. 79:589-620.
- [10] Chestnut BA, Chang Q, Price A, Lesuisse C, Wong M, Martin LJ. 2011. Epigenetic regulation of motor neuron cell death through DNA methylation. *Journal of Neuroscience*. 46:16619-16636.
- [11] Compston A, Coles A. 2002. Multiple sclerosis. *The Lancet*. 359:1221-1231.
- [12] Compston A, Coles A. 2008. Multiple sclerosis. *The Lancet*. 372:1502-1517.
- [13] Copray S, Huynh JL, Sher F, Casaccia-Bonnel P, Boddeke E. 2009. Epigenetic mechanisms facilitating oligodendrocyte development, maturation and aging. *Glia*. 57:5179-1587.
- [14] Dieker J, Muller S. 2010. Epigenetic histone code and autoimmunity. *Clinical Reviews in Allergy and Immunology*. 39:78-84.
- [15] Du C, Liu C, Kang J, Zhao G, Ye Z, Huang S, Li Z, Wu Z, Pei G. 2009. MicroRNA miR-326 regulates TH-17 differentiation and is associated with the pathogenesis of multiple sclerosis. *Nature Immunology*. 10:1252-1259.
- [16] Ebers GC, Yee IM, Sadovnick AD, Duquette P. 2000. Conjugal multiple sclerosis: population-based prevalence and recurrence risks in offspring. *Annals of Neurology*. 48:927-931.
- [17] Ebers GC, Sadovnick AD, Dymont DA, Yee IM, Willer CJ, Risch N. 2004. Parent-of-origin effect in multiple sclerosis: observations in half-siblings. *The Lancet*. 363:1773-1774.
- [18] Feinberg AP, Tycko B. 2004. The history of cancer epigenetics. *Nature Reviews Cancer*. 4:143-153.
- [19] Han L, Witmer PD, Casey E, Valle D, Sukumar S. 2007. DNA methylation regulates microRNA expression. *Cancer Biology and Therapy*. 6:1284-1288.
- [20] Hande, AE, Williamson AJ, Disanto G, Handunnetthi L, Giovannoni G, Ramagopalan SV. 2010. An updated meta-analysis of risk of multiple sclerosis following infectious mononucleosis. *Public Library of Science (PLoS) ONE*. 5:e12496.
- [21] Hernán MA, Jick SS, Logroscino G, Olek, MJ, Ascherio A, Jick H. 2005. Cigarette smoking and the progression of multiple sclerosis. *Brain*. 128:1461-1465.
- [22] Hoppenbrouwers IA, Aulchenko YS, Janssens AC, Ramagopalan SV, Broer L, Kayser M, Ebers GC, Oostra BA, van Duijn CM, Hintzen RQ. 2009. Replication of CD58 and CLEC16A as genome-wide significant risk genes for multiple sclerosis. *Journal of Human Genetics*. 54:676-680.
- [23] Huynh JL, Casaccia P. 2010. Defining the chromatin landscape in demyelinating disorders. *Neurobiology of Disease*. 39:47-52.
- [24] Huynh JL, Casaccia P. 2013. Epigenetic mechanisms in multiple sclerosis: implications for pathogenesis and treatment. *The Lancet Neurology*. 12:195-206.
- [25] [IMSGC] International Multiple Sclerosis Genetics Consortium. 2007. Risk alleles for multiple sclerosis identified by a genome-wide study. *New England Journal of Medicine*. 357: 851-862.

- [26] [IMSGC] International Multiple Sclerosis Genetics Consortium. 2008. Refining genetics associations in multiple sclerosis. *The Lancet Neurology*. 7:567-569.
- [27] Jaenisch R, Bird A. 2003. Epigenetic regulation of gene expression: how the genome integrates intrinsic and environmental stimuli. *Nature Genetics*. 33:245-254.
- [28] Junker A, Krumbholz M, Eisele S, Mohan H, Augstein F, Bittner R, Lassmann H, Werkele H, Hohlfeld R, Meinl E. 2009. MicroRNA profiling of multiple sclerosis lesions identifies modulators of the regulatory protein CD47. *Brain*. 132:3342-3352.
- [29] Koch MW, Metz LM, Kovalchuk O. 2013. Epigenetics and miRNAs in the diagnosis and treatment of multiple sclerosis. *Trends in Molecular Medicine*. 19:23-30.
- [30] Koch MW, Metz LM, Kovalchuk O. 2013. Epigenetic changes in patients with multiple sclerosis. *Nature Reviews Neurology*. 9:35-43.
- [31] Klose RJ, Bird A. 2006. Genomic DNA methylation: the mark and its mediators. *Trends in Biochemical Sciences*. 31:89-97.
- [32] Kürtüncü M, Tüzün E. 2008. Multiple sclerosis: could it be an epigenetic disease? *Medical Hypotheses*. 71:945-947.
- [33] Liggett T, Melnikov A, Tilwalli S, Yi Q, Chen H, Replogle C, Feng X, Reder A, Stefoski D, Balabanov R, et al. 2010. Methylation patterns of cell-free plasma DNA in relapsing-remitting multiple sclerosis. *Journal of Neurological Science*. 290:16-21.
- [34] Liu J, Sandoval J, Doh ST, Cai L, Lopez-Rodas G, Cassacia P. 2010. Epigenetic modifiers are necessary but not sufficient for reprogramming non-myelinating cells into myelin gene-expressing cells. *Public Library of Science (PLOS) ONE*. 5: e13023.
- [35] Mastronardi FG, Wood DD, Mei J, Raijmakers R, Tseveleki V, Dosch HM, Probert L, Cassacia-Bonnefil P, Moscarello MA. 2006. Increased citrullination of histone H3 in multiple sclerosis brain and animal models of demyelination: a role for tumour necrosis factor-induced peptidylarginine deiminase 4 translocation. *Journal of Neuroscience*. 26:11387-11396.
- [36] Mastronardi FG, Noor A, Wood DD, Paton T, Moscarello MA. 2007. Peptidylarginine deiminase 2 CpG island in multiple sclerosis white matter is hypomethylated. *Journal of Neuroscience Research*. 85:2006-2016.
- [37] Milo R, Kahana E. 2010. Multiple sclerosis: geoeidemiology, genetics and the environment. *Autoimmunity Reviews*. 9:387-394.
- [38] Moscarello MA, Wood DD, Ackerly C, Boulias C. 1994. Myelin in multiple sclerosis is developmentally immature. *Journal of Clinical Investigation*. 94:146-154.
- [39] Munger KL, Levin LI, Hollis BW, Howard NS, Ascherio A. 2006. Serum 25-hydroxyvitamin D levels and risk of multiple sclerosis. *Journal of the American Medical Association*. 296:2832-2838.
- [40] Musse AA, Harauz G. 2007. Molecular 'negativity' may underlie multiple sclerosis (MS): role of the myelin basic protein family in the pathogenesis of MS. *International Review of Neurobiology*. 79C: 149-172.
- [41] Nakahara J, Maeda M, Aiso S, Suzuki N. 2012. Current concepts in multiple sclerosis: autoimmunity versus oligodendroglialopathy. *Clinical Reviews in Allergy and Immunology*. 42:26-34.
- [42] Natajara C, Bright JJ. 2002. Curcumin inhibits experimental autoimmune encephalomyelitis by blocking IL-12 signalling through Janus kinase-STAT pathway in T lymphocytes. *Journal of Immunology*. 168: 6506-6513.
- [43] Niller HH, Wolf H, Minarovits J. 2009. Epigenetic dysregulation of the host cell genome in Epstein-Barr virus-associated neoplasia. *Seminars in Cancer Biology*. 19:158-164.
- [44] Nischwitz S, Müller-Myhsok B, Weber F. 2011. Risk conferring genes in multiple sclerosis. *FEBS Letters*. 585:3789-3797.
- [45] Noorbakhsh F, Ellestad KK, Maingat F, Warren KG, Han MH, Steinman L, Baker GB, Power C. 2011. Impaired neurosteroid synthesis in multiple sclerosis. *Brain*. 134: 2703-2721.
- [46] O'Connell RM, Kahn D, Gibson WSJ, Round JL, Scholz RL, Chaudhuri AA, Kahn ME, Rao DS, Baltimore D. 2010. MicroRNA-155 promotes autoimmune inflammation by enhancing inflammatory T cell development. *Immunity*. 33: 607-619.
- [47] Oksenberg JR, Barcellos LF. 2005. Multiple sclerosis genetics: leaving no stone unturned. *Genes and Immunity*. 6:375-387.
- [48] Pedre X, Mastronardi F, Bruck W, Lopez-Rodas G, Kuhlmann T, Casaccia P. 2011. Changed histone acetylation patterns in normal-appearing white matter and early multiple sclerosis lesions. *Journal of Neuroscience*. 31:3435-3445.
- [49] Pereira F, Barbachano A, Silva J, Bonilla F, Campbell MJ, Munoz A, Larriba MJ. 2011. KDM6B/JMJD3 histone demethylase is induced by vitamin D and modulates its effects in colon cancer cells. *Human Molecular Genetics*. 20: 4655-4665.
- [50] Poser CM. 1994. The dissemination of multiple sclerosis: a Viking saga? A historical essay. *Annals of Neurology*. 36:231-234.
- [51] Ramagopalan SV, Ebers GC. 2010. Sequencing the multiple sclerosis genome: more questions than answers. *The Lancet Neurology*. 9:757-759.
- [52] Sadovnick AD. 2012. Genetic background of multiple sclerosis. *Autoimmunity Reviews*. 11:163-166.
- [53] Sevignani C, Calin GA, Siracusa LD, Croce CM. 2006. Mammalian microRNAs: a small world for fine-tuning gene expression. *Mammalian Genome*. 17:189-202.
- [54] Shen S, Casaccia-Bonnefil P. 2008. Post-translational modifications of nucleosomal histones in oligodendrocyte lineage cells in development and disease. *Journal of Molecular Neuroscience*. 35:13-22.
- [55] Smolders J, Menheer P, Kessels A, Damoiseaux J, Hupperts R. 2008. Association of vitamin D metabolite levels with relapse rate and disability in multiple sclerosis. *Multiple Sclerosis Journal*. 14:1220-1224.
- [56] Tsai CN, Tsai CL, Tse KP, Chang HY, Chang YS. 2002. The Epstein-Barr virus oncogene product, latent membrane protein 1, induces the down-regulation of E-cadherin gene expression via activation of DNA methyltransferases. *Proceedings of the National Academy of Sciences*. 99:10084-10089.
- [57] Tullman MJ. 2013. Overview of the epidemiology, diagnosis and disease progression associated with multiple sclerosis. *American Journal of Managed Care*. 19:15-20.
- [58] Wan ES, Weiliang Q, Baccarelli A, Carey VJ, Bacherman H, Rennard SI, Agusti A, Anderson W, Lomas DA, DeMeo DL. 2012. Cigarette smoking behaviours and time since quitting are associated with differential DNA methylation across the human genome. *Human Molecular Genetics*. 21:3073-3082.
- [59] Wei Q, Miskimins WK, Miskimins R. 2005. Stage-specific expression of myelin basic protein in oligodendrocytes involves Nkx2.2-mediated repression that is relieved by the Spl transcription factor. *Advances in Experimental Medicine and Biology*. 280:16284-16294.
- [60] Willer CJ, Dymant DA, Sadovnick AD, Murray TJ, Ebers GC. 2005. Timing of birth and risk of multiple sclerosis: population based study. *British Medical Journal*. 330:120.
- [61] Zhao Q, Rank G, Tan YT, Li H, Moritz RL, Simpson RJ, Cerruti L, Curtis DJ, Patel DJ, Allis CD, et al. 2009. PRMT5-mediated methylation of histone H4R3 recruits DNMT3A, coupling histone and DNA methylation in gene silencing. *Nature Structural and Molecular Biology*. 16:304-311.
- [62] Zhou Y, Simpson RJ, Holloway AF, Charlesworth J, van der Mei I, Taylor BV. 2014. The potential role of epigenetic modifications in the heritability of multiple sclerosis. *Multiple Sclerosis Journal*. 20:135-140.
- [63] Bartel, D.P. 2004. MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell* 116:281-297.
- [64] Mastronardi F. G., Ackerley C. A., Arsenault L, Roots B. I., Moscarello M. A., 1993. Demyelination in a transgenic mouse: A model for multiple sclerosis *Journal of Neuroscience Research* 36 (3): 315-324.