



Bayesian Network based Decision-Support for Leaf Minor Pest Management in Tomato

Niranjan Singh, Neha Gupta

Faculty of Computer Application,

Manav Rachna International University, Faridabad-121004, Haryana, India.

Abstract: Leaf minor (*Liriomyza trifolii*) is one of the key pests of tomato, a prominent vegetable crop grown in northern plains of India. Timely availability of decision-support to the farmers on 'whether and what management option is required' is imperative for effective management of insect pest. For decades, the pest economic threshold level (ETL) has been the basis to select an appropriate pest management option. This process requires quantitative information about pest incidence which needs to be scientifically observed in the farmers' fields. However, a large section of the farming community is not able to scientifically obtain this kind of information. Moreover, in current pest management, decision-making depends upon a large range of agro-ecological information, beside pest incidence. In this study, a Bayesian network-based method/model was devised for selecting appropriate management option of tomato leaf minor on the basis of tentative agro-ecological information that farmers provided. Thus, the resulting method can be used in decision support systems of agriculture with applies information and communication technology to automate and speed up the process of providing pest management decision-support to the farmers.

Keywords: Leaf minor (*Liriomyza trifolii*), Tomato, Decision-Support, ETL, Pest Management

I. Introduction

Tomato (*Lycopersicon esculentum* L.) is an important vegetable crop grown in the northern plains of India [1]. It is grown in 0.458 M ha area with 7.277 M mt production and 15.9 mt/ha productivity in the country. Tomato production and productivity is severely hampered by several insect pests infesting at different stages of the crop growth. The key insect-pests of tomato include thrips (*Thrips tabaci*, *Frankliniella* sp.), aphid (*Myzus persicae*, *Aphis gossypii*), white fly (*Bemisia tabaci*), tobacco caterpillar (*Spodoptera litura*), fruit borer (*Helicoverpa armigera*) and leaf minor (*Liriomyza trifolii*) occurring at different phenological stages of the crop [14]. In spite of regular spraying of insecticides, leaf minor causes considerable crop loss up to 35% [6]. Decision-making is the core of effective pest management [11]. Whenever a farmer notices any pest incidence in his/her field, he/she has to decide 'whether and what pest management action is required', based on the level of pest incidence. The farmers' major concern regarding pest management is timely availability of decision-support on 'whether and what pest management action is required'. In the absence of knowledge and expertise, farmers are over dependent on pesticide dealers for pest management decision-support in the country, which results in excessive, injudicious, and irrational use of chemicals for the pest control. This not only degrades the environment, but also affects human health due to the retention of pesticide residues in vegetables [17]. Presently, agricultural decision support systems (DSSs) available in the country which use information and communication technology (ICT) provide pest management decision-support to the farmers on the basis of the pest economic threshold level (ETL), by collecting scientifically-observed quantitative information about pest incidence from farmers' fields [15]. Nevertheless, a large section of the farming community is neither qualified nor trained to obtain this kind of information. Hence, in the study, a Bayesian network (BN) based method/model has been developed to overcome this problem of decision-support to the farmers. The method could be used in ICT-based agricultural DSSs for the selection of an appropriate pest management option, on the basis of tentative information about agro-ecological factors affecting the pest incidence. The BN approach has been effectively used in ecological decision-making, but only few experiences of its usage have been conducted in pest management [9].

II. Material and Methods

Variety of agro-ecological factors affect pest incidence in the field, but domain experts and earlier studies have reported the presence of natural enemies and unfavourable weather conditions as the most important. The conditions of the crop, along with the level of pest incidence, determine whether and what pest management option needs to be applied [3]. Therefore, meteorological standard week (MSW) wise data records of tomato leaf minor incidence, weather features i.e. maximum temperature and the population of natural enemies (spiders only) for 2013-14, 2014-15 and 2015-16 were obtained from the government institution working in the field of plant protection. These data records were collected from ten farms belonging to major tomato growing villages of Patiala district.

Geographically, Patiala is located at 76.3°E longitude and 30.3°N latitude in the state of Punjab, India. Weekly observations of pest incidence and natural enemies were recorded from two fixed farmer fields (each near to one acre) of tomato crop selected from each village during crop season (October to March). In each farm, five spots were selected randomly, four in the corners at least 5 feet inside of the field borders and one in the centre of the field. Five random plants from each spot were chosen for recording observations. Leaf minor incidence was recorded in terms of no of mines/5leaves/plant. Studies have reported a variety of natural enemies attacking leaf minor at different stages of its lifecycle, but spiders are one of the most important predatory natural enemies of leaf minor [12] which were found in the region. Population of spiders was recorded on a whole plant basis. However, in the region data records revealed very scarce presence of spiders that could have had negligible effect on the pest incidence. This may be due to the indiscriminate use of chemical pesticides by the farmers. Hence, the maximum temperature was the only significant factor which was found affecting the pest incidence. This result was further ascertained through the correlation analysis. The maximum temperature demonstrated a positive role ($r = 0.7738, 0.4268$ and 0.8442) during the years in enhancing the leaf minor fruit infestation. Using these data records as well as the expert elicitation, the BN-based model for decision-making about the selection of an appropriate insect-pests management option was constructed.

Bayesian Network

BNs emerged from research into artificial intelligence, where they were originally developed as a formal means to analyse decision strategies under uncertain conditions [18]. BNs are probabilistic graphical models and consist of 'nodes', that represent a set of random variables, and 'links', that represent their conditional interdependencies (cause-effect relationship) via a directed acyclic graph (DAG) [2], [10]. The values of the nodes are defined in terms of different and mutually exclusive 'states' [8]. A BN can include different types of nodes: 'nature' nodes, 'decision' nodes, and 'utility' nodes. Nature nodes are variables that can be controlled by the decision-maker's actions and represent the empirical or calculated parameters and the probabilities that various states will occur. Input nodes (i.e., nodes without parents) can either be structured as constants or categorical states with associated marginal probability distributions. A decision node represents control variables or events that can directly be implemented by the decision maker. These nodes typically represent the suite of available management actions. Decision nodes should always be accompanied by utility nodes. These utility nodes represent the value of the decisions or outcomes. A utility node can be linked directly to the decision node [7]. Decision nodes do not have probabilities associated with them. BNs that represent and solve decision problems under uncertainty are also known as Bayesian decision networks [16]. As BNs are causal, they can also be used to calculate the effectiveness of interventions, such as alternative management decisions or policies.

A BN is a pair $(G; P)$ [4], [5], [10] where

- $G = (V; E)$ is a DAG whose set of nodes $V = \{X_1, X_2, \dots, X_n\}$ represents the system variables and set of arcs E represents direct dependencies among the variables;
- P is a set of conditional probability distributions containing a conditional probability distribution $P(X_i/pa(X_i))$ for each variable X given the set of parents, $pa(X_i)$ in the graph.

The joint probability distribution over V can be recovered from this set P of conditional probability distributions applying the chain rule as:


$$P(X_1, X_2, \dots, X_n) = \prod_{i=1}^n P(X_i/pa(X_i)) \quad (1)$$

The process of obtaining the probabilities of a BN can be done either manually, from experts' knowledge on the domain, or automatically, from databases [3].

III. Results

On the basis of the relationship between the key factors influencing leaf minor pest incidence and its management options, a BN structure was established (Figure 1) wherein pest incidence, weather (i.e., maximum temperature, presence of spiders, and crop condition were identified as nature nodes, and pest management option was identified as decision node. The arcs indicated the general direction of cause-effect relationship.

Figure 1. BN structure for pest management




Based on the analysis of data records, on expert consultations, and on the numerical definitions that Table 1 provides, all the nodes or variables of the model were discretized.

Table 1. Description of the nodes' states in the Bayesian Network

Node name	Definition	State
Weather	Standard weekly maximum temperature.	Warm: Greater than or equal to 22 OC. Cold: Less than 22 OC.
Presence of natural enemies	Plant wise weekly population of spiders.	Moderate: Greater than or equal to 2.0/plant. Low: Less than 2.0/plant.
Pest incidence	Mines caused by L. minor on plant leaves	High: ≥ 3 mines/5 leaves/plant. Medium: ≥ 1 and < 3 mines/5 leaves/plant. Low: < 1 mines/5 leaves/plant.
Crop condition	Crop health in the field.	Good & Poor (randomized).

The BN model for leaf minor was built using the Netica software (Figure 2). The relationships in the BN are conditionally probabilistic rather than deterministic, except for the utility node.

Figure 2. Bayesian decision network for selection of leaf minor pest management option



Evaluation of model

The evaluation of the model helps to ensure the accuracy and feasibility of the outcome. The model should be evaluated according to the purpose and objective of the exercise. The accuracy of the model should be tested with empirical data, but in many cases these data are not available, at least not voluminously. If data on the system are limited or unavailable, qualitative forms of model evaluation, such as peer review, are valuable [13].

The performance accuracy of the model was measured with leaf minor incidence and weather data records of the selected region available for year 2016-17. Maximum temperature was found as the variable having the most influence on the pest incidence that is an important variable of the model for the selection of a pest management option. The behaviour of pest incidence was tested by applying input values of temperature whereas the value of the presence of natural enemies remained scarce. The software generated probabilities for level of pest incidence which were compared with its true values; the outcomes were found reasonable and logical. The model produced 50% of accuracy. Thus, the pest incidence level probabilities which were generated by the model and the crop condition randomized probabilities were propagated through the model to select an appropriate management option. The BN was also qualitatively evaluated in a meeting involving domain experts, where the model and its results were presented and discussed. The reviewers were satisfied with the results produced by the model as it was intended.

IV. 4. Conclusion

In brief, the BN results were very near to satisfactory. The BN is supposed to be a part of an ongoing iterative model development process, which will be advanced as more knowledge and data relevant to tomato fruit borer incidence are available. The BN which was presented would be considered an initial level of the model which would be further updated. As a result, the final model, which will be developed and validated, can be used in ICT-based agricultural systems to provide real time pest management decisions-support to farmers.

V. Acknowledgements

We are grateful to the ICAR-National Centre for Integrated Pest Management (NCIPM), Government of India for providing pest relevant pest data records.

VI. References

- [1]. S. Bhagat, A. K. Tripathi, I. Ahmad, A. Birah, O. P. Sharma and N. Singh, "Integrated Disease Management for Tomato in Island Ecosystem of Andaman", Indian Journal of Horticulture, vol. 72(1), pp. 67-72, 2015.
- [2]. B. Chunguang and C. Guifen, "Bayesian Networks Modeling for Crop Diseases", In: Li D., Liu Y., Chen Y. (eds) Computer and Computing

- Technologies in Agriculture IV, IFIP Advances in Information and Communication Technology, vol 344. Springer, Berlin, Heidelberg, 2011.
- [3]. M. del Águila and J. del Sagrado, "Metamodeling of Bayesian networks for decision-support system development", In: Proceedings of 8th workshop on knowledge engineering and software engineering (KESE8), 2012.
- [4]. F. V. Jensen and T. D. Nielsen, "Bayesian Networks and Decision Graphs 2nd edition", New York, NY, Springer-Verlag, 2007.
- [5]. U. B. Kjærulff and A. L. Madsen, "Bayesian Networks and Influence Diagrams", New York, NY: Springer-Verlag, 2008.
- [6]. N. K. Krishnakumar, "Bio-ecology and the management of the Serpentine leaf miner, *Liriomyza trifolii* on vegetable crops", Final report of ICAR AD-hoc scheme, IIHR, Bangalore, 1998.
- [7]. M. E. Kragt, "A Beginners Guide to Bayesian Network Modelling for Integrated Catchment Management (Landscape Logic Technical Report No. 9)", Department of Environment, Water Heritage and the Arts, Australia, 2009.
- [8]. R. K. McCann, B. G. Marcot and R. Ellis, "Bayesian Belief Networks: Applications in Ecology and Natural Resource Management", Canadian Journal of Forest Research, vol. 36(12), pp. 3053-3062, 2006.
- [9]. M. Meysam and K. S. Mahdi, "Bayesian Network and Pest Management: A Case Study of Trap Plants on Locust Population", Advances in Environmental Biology, vol. 4(2), pp. 147-151, 2010.
- [10]. J. Pearl, "Probabilistic Reasoning in Intelligent Systems: Networks of Plausible Inference", San Francisco, CA, Morgan Kaufmann Publishers Inc, 1988.
- [11]. V. Rossii, T. Caff and F. Salinari, "Helping Farmers Face the Increasing Complexity of Decision-Making for Crop Protection", Phytopathologia Mediterranea, vol. 51(3), pp. 457-479, 2012.
- [12]. K. Satyagopal, S. N. Sushil, P. Jeyakumar, G. Shankar, O. P. Sharma, S. K. Sain.... S. Saha, "AESA Based IPM Package for Watermelon" Hyderabad, AP, National Institute of Plant Health Management, 2014.
- [13]. H. C. Serena and C. A. Pollino, "Good Practice in Bayesian Network Modelling. Environmental Modelling & Software, vol. 37, pp. 134-145, 2012.
- [14]. K. Singh, "Seasonal Abundance of Fruit Borer, *Helicoverpa armigera* and Its Impact on Marketable Fruit Production in Tomato, *Lycopersicon esculentum* (Mill.)", Agriculture Science Digest, vol. 33(4), pp. 247-252, 2013.
- [15]. N. Singh and N. Gupta, "ICT Based Decision Support Systems for Integrated Pest Management (IPM) in India: A review", Agricultural Reviews, vol. 37(4), pp. 309-316, 2016.
- [16]. N. Singh and N. Gupta, "Decision Making In Integrated Pest Management and Bayesian Network", International Journal of Computer Science & Information Technology, vol. 9(2), pp. 31-36, 2016.
- [17]. N. Singh, H. R. Sardana and S. S. Kumar, "Pest Management Information System (PMIS) for Tomato: A Database Tool for IPM", Indian Journal of Entomology, vol. 74(2), pp. 202-204, 2012.
- [18]. O. Varis, "Bayesian Decision Analysis for Environmental and Resource Management", Environmental Modelling and Software, Vol. 12, pp. 177-185, 1997.