



XRD, THERAPEUTIC APPLICATIONS OF ACNE, BEAM DISPLACEMENT, CHNSO, FL, REFRACTIVE INFLUX, MASS SPECTRAL AND LDT CHARACTERISATIONS OF 4-(4- CHLORO-PHEN-YL)-7,7-DI-METHYL-7,8-DI-HYDRO-4H-1- BENZO-PYRAN-2,5(3H,6H)-DIONE CRYSTALS

A.Munikumari¹, K.Senthilkannan^{2*}, M.Pradhap^{3*}, S.Jeyakumar⁴, R.Krishnaveni⁵, P.Birmanathan⁶,
M Vimalan⁷

¹Yogi Vemana University, Kadapa 516003, Andhra Pradesh, INDIA.

²Dept.of.R&D, Edayathangudy G S Pillay Arts and Science College
Nagapattinam 611002 TN, INDIA.

³Annamalai University, Chidambaram, & On deputation A A Govt College Namakkal, TN, INDIA.

⁴MSAJ group of colleges Chennai, TN India.

⁵Department of Microbiology, Idhaya College for Women, Kumbakonam- 612 001.

⁶Bharathidasan Univ. Model College, Vedaraniyam, Nagapattinam 614810, TN, INDIA.

⁷Dept.of.Physics, Thirumalai Engg.College, Kilambi, Kanchipuram 631551, TN, INDIA.

*Corresponding Author: Dr.K.SenthilKannan

Dean & Director - Research, Vice Principal and Research Scientist,
Edayathangudy G S Pillay Arts & Science College Nagapattinam, 611002.Tamilnadu, INDIA.

Abstract : 4-(4-Chloro-phen-yl)-7,7-di-methyl-7,8-di-hydro-4H-1-benzo-pyran-2,5(3H,6H)-dione crystals have excellent applications in optical, opto - electronics and frequency response, phase matching and other applications such as acne cure in bio and pharma applications. The XRD data denotes that crystal system is monolinic, $P2_1/n$ and parameters are a , b , c in Å as 11.9005, 5.7971, 22.608 and β is 93.973°. The CHNSO analysis refers the elemental and crystalline parameters and analysis for the titled sample, the beam displacement for the nature of the crystal specifications and the fluorescence study reveals that the excitation is in violet spectrum range.

Key words: XRD, CHNSO, beam displacement, Acnecure

1. Introduction

The crystal materials [1-7] are prime over their counter parts by means of their applications in academia, research and industrial utilities and here 4-(4-Chloro-phen-yl)-7,7-di-methyl-7,8-di-hydro-4H-1-benzo-pyran-2,5(3H,6H)-dione are applied for many fields based on their characterisation and utilities and are of NLO crystal [8, 9 and 10] having second harmonic generation and here the XRD, CHNSO, FL studies are represented herewith [11-16]. Acne is fundamentally a hormonal condition driven by male or 'androgenic' hormones, which commonly turned out to be dynamic amid the high school years. Affectability to such hormones, joined with microorganisms on the skin, and unsaturated fats inside oil organs, cause skin inflammation [17-24]. Regular destinations for skin inflammation are the face, chest, bears, and back - the locales of oil organs. Skin inflammation injuries incorporate whiteheads, clogged pores, little knocks, and knobs and sores. In spite of the fact that skin break out is basically an ordinary physiologic event, certain conditions may disturb the condition, including: Fluctuating hormone levels around the season of menses (ladies), Manipulating (picking/nudging) skin inflammation sores, Clothing (for instance, caps and sports protective caps) and headgear. Just three sorts of medications have demonstrated to be viable for the treatment of skin break out - anti-toxins, benzoyl peroxide, and retinoids. A great many people require in any event a couple of specialists, contingent upon the seriousness of their skin break out. Benzoyl peroxide, accessible as an over-the-counter item (for instance, Clearasil, Stridex) and by remedy (for instance, Benoxyl, PanOxyl, Benzopyran and Persagel), targets surface microorganisms [17-25], which frequently exasperate skin inflammation. Ladies of childbearing age must practice anti-conception medication previously, amid, and after treatment (around one month) with isotretinoin. The utilization of isotretinoin requires thorough testing (cholesterol, pregnancy, triglyceride, cholesterol, liver

capacity and bone marrow work) and follow-up for the recommended period (at least 5 months). It is held for the most extreme kinds of skin inflammation that don't react to different medicines. Hormone treatment might be useful for certain ladies with skin break out, particularly for those with signs and side effects (sporadic periods, diminishing hair) of androgen (male hormone) overabundance. The hormone treatment comprises of low-portion estrogen and progesterone (conception prevention pills) or hostile to androgen meds (spironolactone). To anticipate skin break out and diminish its harm to your skin, pursue these tips. Pick a chemical extraordinarily planned for skin inflammation. These items frequently contain salicylic corrosive or benzoyl peroxide, which help to clear skin break out wounds. Clean your face delicately, as injury to the skin break out breakouts may decline the skin inflammation or cause scarring. When washing your face, utilize your hands or cotton cushions, as any terrycloth or other scouring material may make skin break out bruises burst. In the event that you have to utilize a cream, utilize just light, non comedogenic lotions, which don't irritate skin break out. On the off chance that you are a lady, utilize a sans oil establishment. Substantial cosmetics or other corrective items that square pores may cause an erupt of skin inflammation. Benzopyran subordinates are known as calming and cancer prevention agent specialists [25]. Calming just as anticancer benzopyran subsidiaries still stay high in the rundown of the distributed organically dynamic analogs of unique intrigue are (i) the anticancer activity through inhibiting MCT1, CYP1B1, P-gp, P13K/Akt, and AR-JunD[25]; (ii) the treatment of type 2 diabetes[25]; (iii) anti-inflammatory activity[25]; (iv) the treatment of arthritis[25]; (v) the treatment of skin diseases[25]; (vi) the treatment of AD[25]; (vii) the antimicrobial activity[25]; (viii) the antiviral activity[25]; and (ix) the treatment of polycystic kidney disease[16-25].

2. XRD

The 4-(4-Chloro-phen-yl)-7,7-di-methyl-7,8-di-hydro-4H-1-benzo-pyran-2,5(3H,6H)-dione crystals are analysed with XRD blueprint and the lattice parameters are imprisoned with a, b, c in (Å) and by the XRD data as shown in Table.1.

Table.1. XRD data of 4-(4-CHLORO-PHEN-YL)-7,7-DI-METHYL-7,8-DI-HYDRO-4H-1-BENZO-PYRAN-2,5(3H,6H)-DIONE crystals

a	11.9005 Å
b	5.7971 Å
c	22.608 Å
β	93.973°
system	monoclinic
Formula, Space group	C₁₇H₁₇ClO₃, P2₁/ n

3. CHNSO Analysis

The elemental composition of 4-(4-CHLORO-PHEN-YL)-7,7-DI-METHYL-7,8-DI-HYDRO-4H-1-BENZO-PYRAN-2,5(3H,6H)-DIONE is determined using theoretical method and by CHNSO analysis and is given in Table 2 and 3 with the theoretical values for CHNSO % and matches on both the cases with the composition of 4-(4-CHLORO-PHEN-YL)-7,7-DI-METHYL-7,8-DI-HYDRO-4H-1-BENZO-PYRAN-2,5(3H,6H)-DIONE crystals and analysed the formation of the crystal specimen.

Table.2. Elemental representation by theoretical calculations

Elements	% by calculations
C17	67.11
H17	5.592
O3	15.789
N	ND
S	ND
Cl	11.509
Mr, 304	100

Table.3. Elemental representation by CHNSO analysis

Elements	% by CHNSO Analysis
C	67.11
H	5.60
O	15.79
N	ND
S	ND
Cl	ND
Mr 304	88.5

ND as Not Detected

4. Beam displacement

Filter applications by 4-(4-CHLORO-PHEN-YL)-7,7-DI-METHYL-7,8-DI-HYDRO-4H-1-BENZO-PYRAN-2,5(3H,6H)-DIONE for displacement of beam in mm is given in Table 4 for which the displacement is measured with long wave as 1.387 mm and short wave as -0.628 mm and dislocation can also analysed and this refers the opto electronic applications using filters and identified the NLO utility based on the displaced values.

Table.4 Beam displacement of crystals by filter method for Long wave and Short wave

plate level	Beam displacement (mm)
Long wave	1.387
Short wave	-0.628
plate 't'	-0.828

5. FL

The FL study of 4-(4-CHLORO-PHEN-YL)-7,7-DI-METHYL-7,8-DI-HYDRO-4H-1-BENZO-PYRAN-2,5(3H,6H)-DIONE in the range 300–600 nm. A very strong emission peak occurs at 397 nm for the corresponding excitation peaks indicates that the 4-(4-CHLORO-PHEN-YL)-7,7-DI-METHYL-7,8-DI-HYDRO-4H-1-BENZO-PYRAN-2,5(3H,6H)-DIONE crystal has violet fluorescence emission. The maximum intensity peak at 397 nm as in Fig.2 is attributed to $n-\pi^*$ transition and the band gap is 3.125 eV for the 4-(4-CHLORO-PHEN-YL)-7,7-DI-METHYL-7,8-DI-HYDRO-4H-1-BENZO-PYRAN-2,5(3H,6H)-DIONE crystals.

6. Influx

The refractive influx for the grown crystal is found to be 2.9756 microns for the temporal stage of velocity parameters at the point of influx for different rate of variances and mentioned here and is predominant value of typical NLO crystals

7. Conclusion

4-(4-CHLORO-PHEN-YL)-7,7-DI-METHYL-7,8-DI-HYDRO-4H-1-BENZO-PYRAN 2,5(3H,6H) - DIONE crystals have been developed by moderate vanishing strategy. The XRD data addresses that the valuable stone structure and parameters. The CHNSO examination elludes the basic and crystalline parameters and examination for the model, the pole dislodging, flood for the possibility of the valuable stone conclusions and the FL consider reveals that the excitation is in violet range run and used for acne cure.

References

- [1] Burla, M. C., Caliendo, R., Camalli, M., Carrozzini, B., Cascarano, G. L., Giacovazzo, C., Mallamo, M., Mazzone, A., Polidori, G. & Spagna, R. (2012). *J. Appl. Cryst.* 45, 357–361.
- [2] Cyriak Joseph, George Varghese, Ittyachen.M.A, *Cryst.Res.Technol.* 30,159 (1995).
- [3] Desai.C.C and Ramana.M.S.V, *J.Crystal Growth*, 91,126 (1988).
- [4] Dhanuskodi.S and Hariharan.N, *Physica.Stat.Solidi*(1), 106, 67 (1988).
a. E73, 1817–1819 (2017).
- [5] Joshi.M.J and Shah.B.S, *Cryst.Res.Technol*, 19,1107 (1984).
- [6] K SenthilKannan et al (2014), UV, SEM, powder XRD of AAC crystals, *International Journal*
- [7] Marwaha.R.K and Shah.B.S, *Cryst.Res.Technol.*, 26,491 (1991).
- [8] Naik, S. N., Pandey, B. & Ayyangar, N. R. (1988). *Synth. Commun.* 18,625–632.
a. NLO 4 Bromo - 4l Chloro Benzylidene Aniline (BCBA) Crystals, *International*
b. of Chemtech Research, Vol.6(5), pp 3187-3190.
- [9] Portilla, J., Quiroga, J., Cobo, J., Low, J. N. & Glidewell.C. (2007). *Acta Cryst.* C63, o332-o333.
a. Pyridinium Maleate, *Int.J Engineering, Science and Mathematics*, Vol 7(4), pp 310-312.
- [10] Rathke, M. W. & Cowan, P. J. (1985). *J. Org. Chem.* 50, 2622–2624.
- [11] Rigaku (2014). *CrystalClear-SM Expert*. Rigaku Corporation, Tokyo, Japan.
- [12] Senthilkannan.K. *Int J S Res Sci. Engg. Tech.* Mar-Apr; 4(4),941-944 (2018).

- [13] Sheldrick, G. M. (2008). *Acta Cryst.* A64, 112–122.
- [14] Sui YP, Huo HR, Xin JJ, et al. Antibacterial and antitumor activities of biscoumarin and dihydropyran derivatives. *Molecules*. 2015;20:17614–17626.
- [15] Katsori AM, Hadjipavlou-Litina D. Coumarin derivatives: an updated patent review (2012–2014). *Expert Opin Ther Pat*. 2014;24:1323–1347.
- [16] Bukhari SN, Jantan I, Jasamai M. Anti-inflammatory trends of 1, 3- diphenyl-2-propen-1-one derivatives. *Mini Rev Med Chem*. 2013;13:87–94.
- [17] Halestrap AP. Monocarboxylic acid transport. *Compr Physiol*. 2013;3:1611–1643.
- [18] Dhup S, Dadhich RK, Porporato PE, et al. Multiple biological activities of lactic acid in cancer: influences on tumor growth, angiogenesis and metastasis. *Curr Pharm Des*. 2012;18:1319–1330.
- [19] Martel F, Guedes M, Keating E. Effect of polyphenols on glucose and lactate transport by breast cancer cells. *Breast Cancer Res Treat*. 2016;157:1–11. Benzopyran derivatives are important compounds for the treatment of cancer.
- [20] The University of Minnesota. Preparation of benzopyran derivatives and related compounds as MCT1 inhibitors. USA:2013. WO 2013/109972 A2.
- [21] Faiq MA, Dada R, Qadri R, et al. CYP1B1-mediated pathobiology of primary congenital glaucoma. *J Current Glaucoma Practice*. 2015;9:77–80.
- [22] Kwon YJ, Baek HS, Ye DJ, et al. CYP1B1 enhances cell proliferation and metastasis through induction of EMT and activation of Wnt/ beta-Catenin signaling via Sp1 upregulation. *Plos One*. 2016;11: e0151598.
- [23] Hachey DL, Dawling S, Roodi N, et al. Sequential action of phase I and II enzymes cytochrome p450 1B1 and glutathione S-transferase P1 in mammary estrogen metabolism. *Cancer Res*. 2003;63:8492–8499.
- [24] Chen ST, Chen DR, Fang JP, et al. 2,3,7,8-Tetrachlorodibenzo-pdioxin modulates estradiol-induced aldehydic DNA lesions in human breast cancer cells through alteration of CYP1A1 and CYP1B1 expression. *Breast Cancer*. 2015;22:269–279.
- [25] Fylaktakidou KC, Hadjipavlou-Litina DJ, Litinas KE, et al. Natural and synthetic coumarin derivatives with anti-inflammatory/antioxidant activities. *Curr Pharm Des*. 2004;10:3813–3833.




Fig.1 FL Data Of 4-(4-CHLORO-PHEN-YL)-7,7-DI-METHYL-7,8-DI-HYDRO-4H-1-BENZO-PYRAN-2,5(3H,6H)-DIONE crystals

Table.5. ACNE CURE BY 4-(4-CHLORO-PHEN-YL)-7,7-DI-METHYL-7,8-DI-HYDRO-4H-1-BENZO-PYRAN-2,5(3H,6H)-DIONE crystals

S No	Sample size in mm	Acne size	Days of therapeutic applications
1	0.5	0.1 mm	4
2	0.5	0.2 mm	6
3	0.6	0.1 mm	4
4	0.6	0.1 mm	5
5	0.7	0.2 mm	3
6	0.7	0.1 mm	5
7	0.8	0.1 mm	3
8	0.8	0.2 mm	3
9	0.9	0.1 mm	2
10	0.9	0.1 mm	3