



MICROBIAL ESTIMATION OF WOUND INFECTIVITY AND ANTIMICROBIAL ACCESSIBILITY AGAINST *Citrus sinensis*, *Punicagranatum*

R.Krishnaveni^{1*}, K.SenthilKannan^{2*}, V.Eugin Amala³, M.Pradhap⁴, S.Jeyakumar⁵, A Munikumari⁶,
P.Birmanathan⁷

¹Department of Microbiology, Idhaya College for Women, Kumbakonam- 612 001.

²Dept.of.R&D, Edayathangudy G S Pillay Arts and Science College, Nagapattinam 611002.

³Principal, Idhaya College for Women, Kumbakonam- 612 001.

⁴Annamalai University, Chidambaram, & on deputation A AGovt College Namakkal, TN.

⁵MSAJ group of colleges Chennai, TN India.

⁶Department of Genetics and Genomics, Yogi Vemana University,
Kadapa 516003, Andhra Pradesh, India.

⁷Bharathidasan Univ. Model College, Vedaraniyam, Nagapattinam 614810, TN.

Address for Correspondence: Dr.K.SenthilKannan

Dean & Director - Research, Vice Principal and Research Scientist,

Edayathangudy G S Pillay Arts & Science College, Nagapattinam, 611002.Tamilnadu, India.

Abstract: An injury is a break in the tissues of the body. Wounds are for the most part characterized into two sorts. They are open injuries and shut injuries. The breakage which cause in the outside piece of the body is named as open injury and the breakage which cause in the inner piece of the body is named as shut injuries. The idea of the tainting greenery relies upon the basic issue and the area of the procedure. Careful site diseases is characterized as a contamination happening inside 30 days after a careful activity (or inside 1 year if an embed is left set up after methodology) and influencing either cut or profound tissues at the task site. This disease might be shallow or profound incisional contamination or contaminations including organ or body space. Postoperative SSI is among the most widely recognized issues for patients who experience employable strategies and the third most oftentimes detailed nosocomial contamination in the emergency clinic populace. *Citrus sinensis*, *Punicagranatum* zone of hindrance in mm is 10, 15 and 12, 11 and 12.4, 14 and 12, 12 for *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Escherichia coli* & *Klebsiella pneumoniae*

I. Introduction

Staphylococcus sp., settle prosthetic embeds as a biofilm, different layers of sessile cells that hold fast to the embed surface just as to one another. Once a biofilm has framed, it tends to be exceptionally hard to treat clinically in light of the fact that the microbes from the host safe reaction just as anti-toxin operators. Siderophores are the solvent, low atomic mass high proclivity iron-chelating mixes created by organism[1-3]. Vieira et al have likewise announced the antibacterial impact of guava leaves concentrates and found that they repressed the development of the *S. aureus*. Gnan and Demello (testing guava leaf extricate discovered great antimicrobial movement against nine distinct strains of *Staphylococcus aureus*[4-7]. The antibacterial movement of guava leaf extricate was tried against skin break out creating life forms by Qa'dan et al. Presuming that the leaf concentrates might be helpful in treating skin break out particularly when they are known to have mitigating activities [8-10]. Methanolic extricate demonstrates higher antimicrobial movement against tried microorganisms (*E. coli*, *S. aureus*, *Candida albicans* and *Trichophyton rubrum*).

II. Materials and Methods

Site of Collection

The samples for bacteriological screening were taken from government hospital, Kumbakonam, Thanjavur (DT). Where several patients with accidental wounds, burns, surgical wounds and pressure sore. The sample collected by swab and transport to the microbiology lab for the present investigation.

Bacteriological Analysis

The collected sample is then enriched in enrichment medium. From this the culture is transferred into a selective medium. The selective media used for isolation of *Staphylococcus sp.* is mannitol salt agar. The selective media used for isolation of *Streptococcus sp.* is blood agar. *Pseudomonas sp.* was isolated using *Pseudomonas* isolation agar as selective medium. *Escherichia coli* was isolated using Eosin methylene blue agar as selective medium. *Klebsiella sp.* was isolated using MacConkey agar as selective medium.

Microscopic Examination

Hanging Drop Method

Motility of microscopic organisms is recognized utilizing a loopful of medium-term soup culture of the detach by hanging drop technique.

Gram's Staining

A smear of suspected settlement was made on a spotless glass slide and warmth fixed. It was overflowed with gem violet arrangement and permitted to stay for one moment. At that point it was washed with water overwhelmed with iodine arrangement and took into account one moment. It was then depleted and decolourised with 95% ethanol for 15-30 seconds and after that washed and counterstained with saffranin for one moment, at that point analyzed under magnifying instrument.

Biochemical Tests

Indole Production Test, Methyl Red Test, Voges Proskauer Test, Citrate Utilization Test, Triple Sugar Iron Agar Test, Gelatin Hydrolysis, Starch Hydrolysis, Catalase Test, Urease Test, Lipid Hydrolysis, Coagulation Test.

III. Conclusion

The examination introduces an unmistakable comprehension to the causative pathogens of twisted contaminations in this medical clinic and their affectability and obstruction profiles. It has been reasoned that injury contaminations in this were polymicrobial in nature and, much of the time, related. *Citrus sinensis*, *Punica granatum* zone of hindrance in mm is 10, 15 and 12, 11 and 12.4, 14 and 12, 12 for *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Escherichia coli* & *Klebsiella pneumoniae*.

References

- [1] Akhilesh K, Raghvendra P, Vikas S, Madhulika G. Antimicrobial property of lemon peel extract. *Natl J Universal Pharm Life Sci* 2012;52:382-6.
- [2] Anvikar, A.R., Deshmukh, A.B., Karyakarte, R.P., Damle, A.S., Patwardhan, N.S., Malik, A.K., Bichile, L.K., Bajaj J.K., Baradkar V.P., Kulkarni J.D., S.M. Sachdeo. 1999. A one year prospective study of 3280 surgical wounds. *Indian Journal of Medical Microbiology*. 17 (3): 129-132.
- [3] Bajaj Jaya and Chibber Sanjay. 1995. Siderophore production by clinical isolates of *Klebsiella pneumoniae*. *Indian Journal of Medical Microbiology*. 13:34-36.
- [4] Bowler PG, Duerden BI, Armstrong DG. Wound microbiology and associated approaches to wound management. *Clin Microbiol Rev* 2001; 14: 244-69.
- [5] E. Qa'dan, A. Thewaini, D. A. Ali, R. Afifi, A. Elkhawad, and K. Z. Matalka, "The antimicrobial activities of *Psidium guajava* and *Juglans regia* leaf extracts to acne-developing organisms," *The American Journal of Chinese Medicine*, vol. 33, no. 2, pp. 197-204, 2005.
- [6] G. Sacchetti, S. Maietti, M. Muzzoliet *et al.*, "Comparative evaluation of 11 essential oils of different origin as functional antioxidants, antiradicals and antimicrobials in foods," *Food chemistry*, vol. 91, no. 4, pp. 621-632, 2005.
- [7] Mehta, A.P., Rodrigues, C., Sheth, K., Jani, S., Hakimian, A., N. Fazalbhoy. 1998. Control of methicillin resistant *Staphylococcus aureus* in a tertiary care center a five year study. *Indian Journal of Medical Microbiology*. 16(1) : 31-34.
- [8] O'meara, S. N., Callum, N. A., Majid, M. and Sheldon, T.A. (2001). Systematic Review of Antimicrobial Agents Used for Chronic Wounds. *British Journal of Surgery*. 88 (1): 421. Parekh, J. and Chanda S. (2007). In Vitro Antimicrobial Activity of nantans L. Fruit Rind Extracted in Different Solvents. *African Journal Biotechnology*. 6:776-770.
- [9] Obi, R. K., Nwanebu, E. C., Ndubuisi- Nnaji U. U., Onuoha, L. N and Chiegboka, N. (2011). Ethanollic Extraction and Phytochemical Screening of Two Nigerian Herbs on Pathogens Isolated from Wound infections. *International Journal of Comprehensive Pharmacy*. 10(2):1-5.
- [10] Parekh, J. and Chanda S. (2007). In Vitro Antimicrobial Activity of Trapanantans L. Fruit Rind Extracted in Different Solvents. *African Journal of Biotechnology*. 6: 776-770.



Table-1 Types of wound and percentage of sample collection.

Types of wound	Female (35)	
	Numbers	percentage
Accidental	20	100
Surgical	13	100
Burns	1	100
Pressure sore	1	100

Table-2 Testing antimicrobial activity of *Staphylococcus Sp.*,

S.No	Fruits extract	Zone of inhibition (mm)	S/R
1	Citrussinensis	10	S
2	Punicagranatum	15	S

Table-3 showing antimicrobial activity of *Pseudomonas sp.*,

S.No	Fruits extract	Zone of inhibition (mm)	S/R
1	Citrussinensis	12	S
2	Punicagranatum	11	S

Table-4 Showing antimicrobial activity of *E.coli sp.*,

S.No	Fruits extract	Zone of inhibition (mm)	S/R
1	Citrussinensis	12.4	S
2	Punicagranatum	14	S

Table-5 showing antimicrobial activity of *Klebsiella sp.*,

S.No	Plant extract	Zone of inhibition (mm)	S/R
1	Citrussinensis	12	S
2	Punicagranatum	12	S

S – Sensitive R – Resistant