



Mathematical models of tumor growth

Iqbal Kaur

Faculty of Mathematics,
Govt College, Bherian Pehowa, Kurukshetra,
Haryana-India

Abstract: In this research, I exhibit deterministic scientific models regularly utilized for portrayal of tumor development. Advancement of a model, in view of some organic suspicion, is represented by one illustration. Portrayed models are tried and contrasted by capacity with depict trial information. I depict basics of numerical displaying of tumor development and tumor-have associations, and abridge a portion of the fundamental and most unmistakable methodologies. The objective of this model is to gauge future tumor development.

Keywords: Ordinary differential equation, partial differential equation, tumor modeling, angiogenesis, mathematical modeling, tumor growth, multicellular tumor spheroids

I. Introduction

The word tumor is acquired from the Latin word crab since malignancies are unpredictably molded, and like a crab, they "snatch on and don't give up." The term disease particularly alludes to another development which can attack encompassing tissues, metastasize (spread to different organs) and which may inevitably prompt to the patient's demise if untreated. The terms tumor and malignancy are some of the time utilized conversely which can misdirect. A tumor is not really a malignancy. The word tumor just alludes to a mass. For instance, a gathering of liquid would meet the meaning of a tumor. A growth is an especially debilitating kind of tumor. It is useful to keep these qualifications clear while talking about a conceivable malignancy conclusion.

neoplasm-	A neoplasm is an abnormal new development of cells. The cells in a neoplasm generally develop more quickly than ordinary cells and will keep on growing if not treated. As they develop, neoplasms can encroach upon and harm nearby structures. The term neoplasm can allude to benevolent (normally treatable) or harmful (carcinogenic) developments.
tumor-	A tumor is an ordinarily utilized, however non-particular, term for a neoplasm. The word tumor basically alludes to a mass. This is a general term that can allude to benevolent (by and large safe) or threatening (dangerous) developments.
benign tumor-	Benign tumors are non-dangerous/non-malignant tumor. An amiable tumor is typically confined, and does not spread to different parts of the body. Most kindhearted tumors react well to treatment. Be that as it may, if left untreated, some amiable tumors can develop huge and prompt to genuine ailment due to their size. Generous tumors can likewise emulate harmful tumors, thus consequently are infrequently treated.
malignant tumor-	Malignant tumors are cancerous growths. They are often resistant to treatment, may spread to other parts of the body and they sometimes recur after they were removed.
cancer-	A cancer is another word for a malignant tumor (a malignant neoplasm).

Cell division is fundamental for solid development of a life form and complex hereditary systems have advanced to switch cell division on and off at the best possible time for ordinary improvement. Disease happens when cells separate wildly and create a developing protuberance or tumor. Tumors give out substance flags that provoke the development of veins to supply them with oxygen and sustenance. An especially risky part of tumor is metastasis. Here, little bits of the tumor sever and make a trip through the circulatory system to attack different parts of the body.

Angiogenesis is the procedure by which tumors instigate veins from the host tissue to grow narrow tips which move towards and at last enter the tumor, furnishing it with a circling blood supply and, along these lines, a practically boundless wellspring of supplements.

Multicellular spheroids (MCS) are groups of disease cells, utilized as a part of the lab to concentrate the early phases of avascular tumor development. Develop MCS have an all around characterized structure, containing a focal center of necrotic, or dead, cells, encompassed by a layer of non-multiplying, tranquil cells, with multiplying cells confined to the external, supplement rich layer of the tumor. Accordingly, they are regularly used to evaluate the adequacy of new hostile to disease medications and treatment treatments.

Numerical models of MCS commonly comprise of a conventional differential condition (ODE) coupled to at least one response dispersion conditions (RDEs). The ODE is gotten from mass protection and depicts the advancement of the external tumor limit, though the RDEs portray the dispersion inside the tumor of fundamental supplements, for example, oxygen and glucose and development inhibitors. These models precisely imitate the development of spheroids furthermore the naturally visible heterogeneity which is the sign of spheroids. The model created in this paper displays comparative elements. Be that as it may it likewise empowers us to survey a tumor's potential for intrusion. The key element that decides this potential is the harmony between an interior far reaching power (brought on by cell expansion) and a controlling power (created by strengths of attachment which exist between cells on the tumor limit).

A mathematical model of tumor growth is a mathematical expression of the dependence of tumor size on time. There are three main steps in the process of mathematical modeling:

1. Definition of model based on biological assumptions.
2. Testing the model against experimental data
3. Acceptance of model or its rejection and change of assumptions.

II. ORDINARY DIFFERENTIAL EQUATION MODELS OF TUMOR GROWTH

let $x(t+\Delta t)$ =no of tumor cells at time $t+\Delta t$
and $x(t)$ =no of tumor cells at time t

then,

$$\frac{dx}{dt} = \lim_{\Delta t \rightarrow 0} \frac{x(t+\Delta t) - x(t)}{\Delta t}$$

$$\frac{dx}{dt} = \lambda x$$

As before, if $\lambda < 0$, $\lambda = 0$, $\lambda > 0$ the population decreases, remains at a constant size, or increases respectively.

However, there are mainly two types of growth models for tumors: exponential and non-exponential. **Exponential growth model** According to the Exponential growth model, tumor volume increases exponentially by time:

$$V = V_0 e^{\alpha t}, \quad (1)$$

where α is the exponential growth constant, and V and V_0 are the tumor volume at times t and 0 , respectively. This model implies that the tumor volume can increase indefinitely and the growth rate of tumor is proportional to its volume

$$\frac{dV}{dt} = \alpha V \quad (2)$$

The simplest theory of tumor growth presumes all cells proliferate with constant cell cycle duration T_c . This leads to exponential growth, which is also valid in the extended cases where either a constant fraction of the volume is proliferating or the cell cycle length is a random variable with exponential distribution (assuming that the individual cell cycle length distributions are independent and identically distributed). As one modification, initial exponential phase can be assumed to be followed by a linear growth phase [39], giving the following Cauchy problem for the volume rate of change (growth rate):

$$\frac{dV}{dt} = \alpha V, \quad t \leq u$$

$$\frac{dV}{dt} = \dot{\alpha}, \quad t > u$$

$$V(t=0) = V_0$$

Here, the coefficient $\dot{\alpha}$ is the fraction of proliferative cells times $\ln 2/T_c$ where T_c is either the constant cell cycle length or the mean cell cycle length (under the assumption of exponentially distributed cell cycle lengths). The coefficient $\dot{\alpha}$ drives the linear phase. Assuming that the solution of the problem (1) is continuously differentiable uniquely determines the value of u as $u = \log_{\frac{\dot{\alpha}}{\alpha V_0}}$. The coefficient denotes the initial volume. From this formula, three models were considered: a) initial volume fixed to 1 mm³ and no linear phase, referred to hereafter as exponential 1, b) free initial volume and no linear phase, referred to as exponential and c) equation (1) with fixed initial volume of mm³, referred to as the exponential-linear model.

Tumor volume doubling time (DT) Tumor volume doubling time, DT, was introduced for quantification of tumor growth rate in 1956 when Collins et al. proposed a graphical method to estimate the DT of tumors (Collins,

Loeffler et al. 1956). DT is the time needed for a tumor to double in volume. The mathematical approach for estimation of DT was then proposed in 1961 (Schwartz 1961):

$$\ln(V_2/V_1) / (t_2 - t_1) = \ln 2 / DT \quad (3)$$

where V_1 and V_2 are two tumor volume estimates at two different occasions, t_1 and t_2 , respectively. DT has been widely used as a quantity for tumor growth rate since its introduction. There are flaws with DT as a quantity for tumor growth rate: the frequency distribution of DT in a population is not normal and there are tumors with very long DT values in a population (Spratt 1969). Therefore, mean DT, DT_m , does not indicate the average growth rate and DT is not suitable for common statistical testing. Some researchers have approximated the frequency distribution of DT by a log-normal distribution (Spratt and Spratt 1964; Shackney, McCormack et al. 1978; Balmukhanov, Turdugulov et al. 1982; Kuroishi, Tominaga et al. 1990; Usuda, Saito et al. 1994). The average growth rate is then estimated by DT_{log} , calculated as the antilog to the arithmetic mean of the logarithms of doubling times (Spratt 1969; Gregory, Richards et al. 1991; Spratt, Meyer et al. 1995). The logarithm of DT, $\log(DT)$, is also proposed to be more suitable for statistical testing (Spratt 1969). DT_{log} is mathematically equal to geometric mean DT, DT_{gm} , which is also used to estimate the average growth rate (Kuroishi, Tominaga et al. 1990; Blomqvist, Wiklund et al. 1993; Usuda, Saito et al. 1994). It is also clear from Eq. 3 that DT is not defined when the estimated tumor volumes are equal. The reason for these flaws with DT has not previously been studied.

Non-exponential tumor growth models Studies have shown that tumor growth rate may decline with time (Hart, Shochat et al. 1998; Bajzer 1999; Afenya and Calderon 2000), which results in non-exponential growth model of tumors. Growth deceleration has been observed in animal models (Wennerberg, Willen et al. 1988), for solid tumors in clinical studies (Spratt, von Fournier et al. 1993; Spratt, Meyer et al. 1996), and in leukemia (Afenya and Calderon 2000). Growth deceleration is attributed to several factors, including prolonged cell cycle, reduced growth fraction, decreased availability of oxygen (Pavelic, Porter et al. 1978), decreased cell proliferation rate with increased cell loss rate (Bassukas and Maurer-Schultze 1987), tumor-related systemic factors (DeWys 1972), and allometric growth control (Prehn 1991). A number of non-exponential growth models are available in the literature, among which the Gompertzian model is widely used (Araujo and McElwain 2004)

III. DISCUSSION

An expanding assortment of numerical models has advanced into growth examine over the recent decades. In this we have delineated how basic quantitative models are produced and contrasted and exploratory information, and demonstrated how they can be utilized to recreate complex organic procedures and communications. We have picked original papers as illustrations, and for effortlessness have needed to forget a vast assemblage of brilliant scientific demonstrating writing. The intrigued per user is alluded to late audit articles and books that give a more comprehensive outline of the condition of-the art in tumor displaying.

References

- [1] L. von Bertalanffy, Untersuchungen über die Gesetzlichkeit des Wachstums, VII, Stoffwechseltypen und Wachstumstypen Biol. Zentralbl. 61(1941), 510–535.
- [2] L. von Bertalanffy, Quantitative laws in metabolism and growth, Q. Rev. Biol. 32(1957), 217–231.
- [3] R. Bjerkvig (Ed.), Spheroid Culture in Cancer Research. CRC Press, Boca Raton, 1992. 190 M. Marušić
- [4] A. D. Conger, M. C. Ziskin, Growth of mammalian multicellular tumor spheroids, Cancer Res. 43(1983), 556–580.
- [5] R. D. Cook, S. Weisberg, Linear and Nonlinear Regression, in: Statistical Methodology in the Pharmacological Sciences, D. A. Berry, ed., Marcel Dekker, New York, 1990, 163–199.
- [6] E. B. Cox, M. A. Woodbury, L. E. Myers, A new model for tumor growth analysis based on a postulated inhibitory substance, Comp. Biomed. Res. 13(1980) 437.
- [7] J. Durbin, G. S. Watson, Testing for serial correlation in least squares regression. I. Biometrika 37(1950), 409–428.
- [8] J. Durbin, G. S. Watson, Testing for serial correlation in least squares regression. II. Biometrika 38(1951), 159–177.
- [9] J. P. Freyer, Role of necrosis in regulating the growth saturation of multicellular spheroids, Cancer Res. 48(1988), 2432–2439.
- [10] J. P. Freyer, R. M. Sutherland, Regulation of growth saturation and development of necrosis by the glucose and oxygen supply Cancer Res. 46(1986), 3513–3520.
- [11] B. Gompertz, On the nature of the function expressive of the law of human mortality, Philos. Trans. Roy. Soc. London 36(1825), 513.
- [12] E. Kreyszig, Introductory Mathematical Statistics, Wiley, New York, 1970.
- [13] J. Landry, J. P. Freyer, R. M. Sutherland, A model for the growth of multicellular spheroids, Cell Tissue Kinet. 15(1982), 585–594.
- [14] S. A. Maggelakis, J. A. Adam, Mathematical model of prevascular growth of a spherical carcinoma, Math. Comput. Modeling 13(1990), 23–38.
- [15] M. Marušić, Z. Bajzer, Generalized two-parameter equation of growth, J. Math. Anal. Appl. 179(1993), 446–462.
- [16] M. Marušić, Z. Bajzer, J. P. Freyer, S. Vuk-Pavlović, Modeling autostimulation of growth in multicellular tumor spheroids, Int. J. Biomed. Comput. 29(1991), 149–158.
- [17] M. Marušić, Z. Bajzer, J. P. Freyer, S. Vuk-Pavlović, Analysis of growth of multicellular tumour spheroids by mathematical models, Cell Prolif. 27(1994), 73–94. Mathematical model of tumor growth 191

- [18] M. Marušić, S. Vuk-Pavlović, Prediction power of mathematical models for tumor growth, *J. of Biological Systems* 1(1993), 69–78. [19] J. J. More: The Levenberg-Marquardt algorithm: Implementation and theory, in: *Numerical Analysis*, G. A. Watson, ed., Springer, New York, 1977.
- [20] R. Pearl: *Studies in Human Biology*, Williams & Wilkins, Baltimore, 1924.
- [21] S. Piantadosi, A model of growth with first-order birth and death rates, *Comp. Biomed. Res.* 18(1985), 220–232.
- [22] W. H. Press, B. P. Flannery, S. A. Teukolsky, W. T. Vetterling, *Numerical Recipes*, Cambridge University Press, Cambridge, 1986.
- [23] M. A. Savageau, Allometric morphogenesis of complex systems: a derivation of the basic equations from first principles, *Proc. Natl. Acad. Sci. USA* 76(1979), 6023–6025.
- [24] G. Schwarz, Estimating the dimension of a model, *Ann. Stat.* 6(1978), 461–464.
- [25] L. F. Shampine, M. K. Gordon, *Computer Solution of Ordinary Differential Equations. The Initial Value Problem*, Freeman, New York, 1975.
- [26] S. Siegel, N. J. Castellan Jr., *Nonparametric Statistics for the Behavioral Sciences*, 2nd edition, McGraw-Hill, New York, 1988.
- [27] R. M. Sutherland, Cell and environment interactions in tumor microregions: the multicell spheroid model, *Science* 240(1988), 177.
- [28] M. E. Turner Jr., E. L. Bradley Jr., K. A. Kirk, K. M. Pruitt, A theory of growth, *Math. Biosci.* 29(1976), 367–373.
- [29] V. G. Vaidya, F. J. Alexandro Jr., Evaluation of some mathematical models for tumor growth, *Int. J. Biomed. Comput.* 13(1982), 19–35.
- [30] P. F. Verhulst, Notice sur la loi que la population suit dans son accroissement, *Corr. Math. Phys.* 10(1838), 113–121.
- [31] T. E. Wheldon, *Mathematical Models in Cancer Research*, Adam Hilger, Bristol, 1988.
- [32] T. E. Wheldon, J. Kirk, W. M. Grey, Mitotic autoregulation, growth control and neoplasia, *J. Theor. Biol.* 38(1973), 627. 192 M. Marušić
- [33] Benzekry Sébastien Clare Lamont, Afshin Beheshti Amanda Tracz, John M. L. Ebos,, Lynn Hlatky, Philip Hahnfeldt, Classical “Mathematical Models for Description and Prediction of Experimental Tumor Growth” computational biology Published: August 28, 2014
- [34] Mehrara Esmaeil Quantitative analysis of tumor growth and response to therapy